Literature Cited

Batiuk, R.A., Orth, R.J., Moore, K.A., Dennison, W.C., Stevenson, J.C., Staver, L.W., Carter, V., Rybicki, N.B., Hickman, R.E., Kollar, S., Bieber, S., Heasly, P. 1992. *Chesapeake Bay Submerged Aquatic Vegetation Habitat Requirements and Restoration Targets: A Technical Synthesis*. Chesapeake Bay Program, Annapolis, MD., CBP/TRS 83/92, Contract No. 68-WO-0043. 248 p.

Benoy, G.A., Kalff, J. 1999. Sediment accumulation and Pb burdens in submerged macrophytee beds. Limnology and Oceanography 44(4) 1081-1090.

Bigley, R.E. 1981. The population biology of two intertidal seagrasses, *Zostera marina* and *Ruppia maritima*, at Roberts Bank, British Columbia. Thesis, University of British Columbia, Vancouver British Columbia, Canada.

Borum, J. 1985. Development of epiphytic communities on eelgrass (*Zostera marina*) along a nutrient gradient in a Danish Estuary. Marine Biology 87:211-218.

Brush, G.S., Davis, F.W. 1984. Stratigraphic evidence of human disturbance in an estuary. Quaternary Research 22:91-108.

Brush, G.S., Hilgartner, W.B. 2000. Paleoecology of submerged macrophytes in the upper Chesapeake Bay. Ecological Monographs 70:645-667.

Burdick, D.M., Kendrick, G.A. 2001. Standards for seagrass collection, identification, and sample design. In: Short, F.T., Coles, R.G., editors. Global Seagrass Research Methods. Amsterdam: Elsevier Science BV. p 79-100.

Burt, J.S., Kendrock, G.A., Mansini, R. J., Simpson, C.J. 1995. Light and *Posidonia sinuosa* seagrass meadows in the temperate coastal waters of Western Australia: II. Effect of epiphyte species assemblage and biomass on attenuating light to the leaf surface.

Technical Series 62. Department of Environmental Protection, Perth, Western Australia.

Cabin, R.J., Marshall, D.L., Mitchell, R.J. 2000. The demographic role of soil seed banks. II. Investigations of the fate of experimental seeds of the desert mustard *Lesquerella fendleri*. Journal of Ecology 88:293-302.

Cambridge, M.L., McComb, A.J. 1984. The loss of seagrasses in Cockburn sound, Western Australia. I. The time course and magnitude of seagrass decline in relation to industrial development. Aquatic Botany 20:229-243.

Chambers, J.C., MacMahon, J.A. 1994. A day in the life of a seed: movements and fates of seeds and their implications for natural and managed systems. Annual Review of Ecology and Systematics 25:263-292.

Cook, R.E. 1979. Patterns of juvenile mortality and recruitment in plants. In: Solibrig, O.T., Jain, S., Johnson, G.B., Raven, P.H., editors. Topics in Plant Population Biology. New York: Columbia University Press. p 207-231.

Den Hartog, C., Polderman, P.J.G. 1975. Changes in seagrass populations of the Dutch Waddenzee. Aquatic Botany 1:141-147.

Dennison, W.C., Albert, R.S. 1986. Photoadaptation and growth of *Zostera marina L*. (eelgrass) transplants along a depth gradient. Journal of Experimental Marine Biology and Ecology 98:265-282.

Dennison, W.C. 1987. Effects of light on seagrass photosynthesis, growth, and depth distribution. Aquatic Botany 27:15-26.

Dennison, W.C., Orth, R.J., Moore, K.A., Stevenson, J.C., Carter, V., Kollar, P.W., Bergstrom, P.W., Batiuk, R.A. 1993. Assessing water quality with submersed aquatic vegetation. BioSience 143:86-94.

Fishman, J.R., Orth, R.J. 1996. Effects of predation on *Zostera marina* seed abundance. Journal of Experimental Marine Biology Ecology 198:11-26.

Fonseca, M.S., Fisher, J.J., Zieman, J.C., Thayer. G.W. 1982. Influence of the seagrass, *Zostera marina* L., on current flow. Estuarine Coastal Shelf Science. 15:351-364.

Fonseca, M.S. 1992. Restoring seagrass in the Inited States. In: Thayer, G.W., editor. Restoring the nation's marine environment, Maryland Sea Grant, College Park, MD. p 79-110.

Fonseca, M.S., Kenworthy, W.J., Courtney, F.X., Hall, M.O. 1994. Seagrass planting in the southeastern United States: Methods for accelerating habitat development.

Restoration Ecology 2(3):198-212.

Fonseca, M.S., Kenworthy, W.J., Thayer, G. 1998. Guidelines for the conservation and restoration of seagrasses in the United States and adjacent waters. In: NOAA Coastal Ocean Office, Silver Spring, MD. 222 p.

Gacia, E., Granata, T.C., Duarte, C.M. 1999. An approach to measurement of particle flux and sediment retention within seagrass meadows. Aquatic Botany 65:255-268.

Granger, S., Traber, M., Nixon, S.W., Keyes, R. 2002. A practical guide for the use of seeds in eelgrass (Zostera marina L.) restoration. Part I. Collection, processing, and storage. Schwartz, M. (editor), Rhode Island Sea Grant, Narragansett, R.I. 20 p.

Harper, J.L., Williams, J.T., Sagar, G.R. 1965. The behavior of seeds in soil. 1. The heterogeneity of soil surfaces and its role in determining the establishment of plants from seed. Journal of Ecology 53:273-286.

Harrison, P.G. 1987. Variations in success of eelgrass transplants over a five-years' period. Environmental Conservation 17(2):157-163.

Harwell, M.C., Orth, R.J. 1999. Eelgrass (*Zostera marina* L.) seed protection for field experiments and implications for large scale restoration. Aquatic Botany 64:51-61.

Inglis, G.J., Waycott, M. 2001. Methods for assessing seagrass ecology and population genetics. In: Short, F.T., Coles, R.G., editors. Global Seagrass Research Methods.

Amsterdam: Elsevier Science BV. p 123-140.

Janzen, D.H. 1971. Seed predation by animals. Annual Review of Ecology and Systematics 2:465-492.

Kemp W.M., Boynton, W.R., Stevenson, J.C., Twilley, R.R., Means, J.C. 1983. The decline of submerged vascular plants in upper Chesapeake Bay: Summary of results concerning possible causes. Marine Technology Society Journal 17:78-89.

Kemp, W. M., W. R. Boynton, A. J. Hermann. 1995. Simulation models of an estuarine macrophyte ecosystem. pp. 262-278, In: Patten, B., Jørgensen, S.E., editors. Complex ecology. Prentice Hall, Englewood Cliffs, NJ.

Luckenbach, M.L., Orth, R.J. 1999. Effects of a deposit-feeding invertebrate on the entrapment of *Zostera marina* L. seeds. Aquatic Botany 62:235-247.

Madden, C., Day, J. 1992. Induced turbulence in rotating bottles affects phytoplankton productivity measurements in turbid waters. Journal of Plankton Research 14(8):1171-1191.

Maryland Department of Natural Resources (MD-DNR). 2005. *Maryland Tributary*Strategy Patuxent River Basin Summary Report for 1985-2003 Data. 61 p. [Electronic version.] Retrieved July 27, 2005 from

http://www.dnr.state.md.us/bay/tribstrat/basin_summary_pax_012505.pdf

McRoy, C.P. 1968. The distribution and biogeography of *Zostera marina* (eelgrass) in Alaska. Pacific Science 22:507-513.

Moore, K.A., Orth, R.J., Nowak, J.F. 1993. Environmental regulation of seed germination in *Zostera marina* L. (eelgrass) in Chesapeake Bay: Effects of light, oxygen, and sediment burial depth. Aquatic Botany 45:79-91.

Moore, K.A., Neckles, H.A., Orth R.J. 1996. *Zostera marina* L. (eelgrass) growth and survival along a gradient of nutrients and turbidity in the lower Chesapeake Bay. Marine Ecology Progress Series 142:247-259.

Moore, K.A., Wetzel, R.L., Orth, R.J. 1997. Seasonal pulses of turbidity and their relations to eelgrass (*Zostera marina* L.) survival in an estuary. Journal of Experimental Marine Biology and Ecology 215:115-134.

Naylor, M.D., Kazyak, P. 1995. Quantitative characterization of submerged aquatic vegetation species in tidal freshwater reaches of the Patuxent River drainage basin.

Submitted to Maryland Department of the Environment. Maryland Department of Natural Resources, Annapolis, MD. 24 p.

Olesen, B., Sand-Jensen, K. 1994. Patch dynamics of eelgrass *Zostera marina*. Marine Ecology Progress Series 106:147-156.

Olesen, B. 1999. Reproduction in Danish eelgrass (*Zostera marina L.*). Marine Ecology Progress Series 106:147-156.

Orth, R.J. 1975. Destruction of eelgrass, *Zostera marina* by the cownose ray, *Rhinoptera bonasus*, in Chesapeake Bay. Chesapeake Science 16:205-208.

Orth, R.J., Moore, K.A. 1983. Chesapeake Bay: An unprecedented decline in submerged aquatic vegetation. Science 222:51-53.

Orth, R.J., Luckenbach, M.W., Moore, K.A. 1994. Seed dispersal in a marine macrophyte: implications for colonization and restoration. Ecology 75:1927-1939.

Orth, R.J., Nowak, J.F., Wilcox, D.J., Whiting, J.R., Nagey, L.S. 1999. Distribution of submerged aquatic vegetation in the Chesapeake Bay and tributaries and Chincoteague Bay – 1998. Final Report, US Environmental Protection Agency, Annapolis, MD.

Orth, R.J., Harwell, M.C., Bailey, E.M., Batholomew, A., Jawad, J., Lombara, A., Moore, K., Rhode, J., Woods, H. 2000. A review of issues in seagrass seed dormancy and germination: Implications for conservation and restoration. Marine Ecology Progress Series 200:277-288.

Orth, R.J., Fishman, J.R., Harwell, M.C., Marion, S.R. 2003. Seed-density effects on germination and initial seedling establishment in eelgrass *Zostera marina* in the Chesapeake Bay region. Marine Ecology Progress Series 250:71-79.

Orth, R.J., Bieri, J., Fishman, J.R., Harwell, M.C., Marion, S.R., Moore, K.A., Nowak, J.F., van Montfrans, J. In Press. A review of techniques using adult plants and seeds to transplant eelgrass (*Zostera marina* L.) in Chesapeake Bay and the Virginia Coastal Bays. Proc. Conf. Seagrass Restoration: Success, Failure, and the Costs of Both. March 11, 2003. Sarasota, Florida.

Parham, T., Karrh, L. 1998. *Maryland Submerged Aquatic Vegetation Restoration Targeting System.* Final Report to National Oceanic and Atmospheric Administration.

Maryland Department of Natural Resources, Annapolis, MD. 32 p.

Patuxent River Commission Staff. 2003. Patuxent River Basin Summary. p 1-2.

Petticrew, E.L., Kalff, J. 1991. Predictions of surficial sediment composition in the littoral zone of lakes. Limnology and. Oceanography. 36:384-392.

Phillips, G.L., Emnison, D., Moss, B. 1978. A mechanism to account for macrophyte decline in progressively eutrophicated freshwaters. Aquatic Botany 4:103-126.

Pickerell, C., Schott, S., Wyllie-Echeverria, S. 2003. Buoy-deployed seeding: A new approach to restoring seagrass using seed. In: Submerged Aquatic Vegetation Propagation Workshop; 2003 September 3-4; Baltimore, MD. http://el.erdc.usace.army.mil/sav/workshop.html

Pickerell, C., Schott, S. and Wyllie-Echeverria, S. 2005. Buoy Deployed Seeding: Demonstration of a new eelgrass (*Zostera marina* L.) planting method. Ecological Engineering 25:127-136.

Rhoads, D.C., Young, D.K. 1970. The influence of deposit feeding organisms on sediment stability and community trophic structure. Journal of Marine Research 28:150-178.

Rhoads, D.C. 1974. Organism-sediment relations on the muddy sea floor. Oceanography and Marine Biology: An Annual Review. 12:263-300.

Silberhorn, G.M., Orth, R.J., Moore, K.A. 1983. Anthesis and seed production in *Zostera marina* L. (eelgrass) from the Chesapeake Bay. Aquatic Biology 15:133-144.

Stankelis, R.M., Boynton, W.R., Frank, J.M. 1999. Submerged aquatic vegetation (SAV) habitat evaluation. Boynton, W.R., Rohland, F.M., editors. Ecosystem Processes

Component Level 1 Interpretive Report No 16. Chesapeake Biological Laboratory,

University of Maryland System. p 77-117.

Stankelis, R.M., Naylor, M.D., Boynton, W.R. 2003. Submerged aquatic vegetation in the mesohaline region of the Patuxent estuary: Past, present, and future status. Estuaries 26:186-195.

Thom, R.M., Borde, A.B., Williams, G.D., Southared, J.A., Blanto, S.L., Woodruff, D.L. 2001. Effects of multiple stressors on eelgrass restoration projects. Proceedings of Puget Sound Research 2001, The fifth Puget Sound Research Conference, 2001.

Twilley, R.R., Kemp, W.M., Staver, K.W., Stevenson, J.C., Boynton, W.R. 1985.

Nutrient enriched of estuarine submerged vascular plant communities. 1. Algal growth and effects on production of plants and associated communities. Marine Ecology Progress Series 23:179-191.

Wassenberg, T.J. 1990. Seasonal feeding on *Zostera capricorni* seeds by juvenile *Peneaus esculentus* (Crustacea: Decapoda) in Moreton Bay, Queensland. Australian Journal of Marine and Freshwater Research 41:301-310.

Wetzel, R.G., and Hough, R.A. 1973. Productivity and role of aquatic macrophytes in lakes: An assessment. Polish Archives of Hydrobiology 20:9-19.

Wetzel, R.G., Penhale, P.A. 1983. Production ecology of seagrass communities in the lower Chesapeake Bay. Marine Technology Society Journal 17:22-31.